

Anomalous transport: Theory and Applications - II

Karl Landsteiner

Instituto de Física Teórica UAM-CSIC



Holography and Topology of Quantum Matter, APCTP, Pohang 2016

$$\vec{J}_A = d_{ABC} \frac{\mu_B}{4\pi^2} \vec{B}_C + \left(d_{ABC} \frac{\mu_B \mu_C}{4\pi^2} + b_A \frac{T^2}{12} \right) \vec{\Omega}$$

$$\vec{J}_\epsilon = \left(d_{ABC} \frac{\mu_B \mu_C}{8\pi^2} + b_A \frac{T^2}{24} \right) \vec{B}_A + \left(d_{ABC} \frac{\mu_A \mu_B \mu_C}{6\pi^2} + b_A \frac{\mu_A T^2}{6} \right) \vec{\Omega}$$

Outline

- Relativistic Hydrodynamics
- Kubo Formulas
- Anomalous Transport from Holography
- Renormalization
- Application: CME in QGP
- Application: NMR in WSM

Relativistic Hydro

- Thermodynamics: maximal entropy = forget everything that can be forgotten (conserved charges)
 - Energy: Temperature T
 - Charge: Chemical potential μ
 - Time: Frame u^μ $u^2 = -1$

$$Z = \text{Tr} e^{-\beta(u^\mu P_\mu + \mu Q)}$$

$$T^{\mu\nu} = (\epsilon + p)u^\mu u^\nu + pg^{\mu\nu}$$

$$J^\mu = \rho u^\mu$$

$$p = p(T, \mu) \quad , \quad \epsilon + p = sT + \rho\mu$$

Relativistic Hydro

- Hydrodynamics: hypothesis of local thermal equilibrium
- Universal low energy, long wavelength behaviour
- Equations of motion: Conservation laws
- Ideal Hydrodynamics: $T(x), \mu(x), u^\mu(x)$

$$\partial_\mu T^{\mu\nu} = 0$$

$$\partial_\mu J^\mu = 0$$

$$s^\mu = s u^\mu$$

$$\partial_\mu s^\mu = 0$$

Relativistic Hydro

- Viscous Hydrodynamics: add derivatives
- Effective Field Theory: all terms consistent with symmetries and 1 derivative

$$T^{\mu\nu} = (\epsilon + p)u^\mu u^\nu - pg^{\mu\nu} + \tau^{\mu\nu}$$

$$J^\mu = \rho u^\mu + \nu^\mu$$

- Ambiguities in local definitions of $T(x)$, $\mu(x)$, $u^\mu(x)$
- Frame choice: no corrections to ϵ, p and $u^\mu \tau^{\mu\nu} = 0$ (Landau frame)

$$\tau^{\mu\nu} = -\eta \mathcal{P}^{\mu\alpha} \mathcal{P}^{\nu\beta} \left(\partial_\alpha u_\beta + \partial_\beta u_\alpha - \frac{2}{3} \partial_\lambda u^\lambda g_{\alpha\beta} \right) - \zeta \mathcal{P}^{\mu\nu} \partial_\lambda u^\lambda$$

$$\nu^\mu = \sigma \left[\mathcal{P}^{\mu\alpha} \partial_\alpha \left(\frac{\mu}{T} \right) - E^\mu \right]$$

Relativistic Hydro

$$\tau^{\mu\nu} = -\eta \mathcal{P}^{\mu\alpha} \mathcal{P}^{\nu\beta} \left(\partial_\alpha u_\beta + \partial_\beta u_\alpha - \frac{2}{3} \partial_\lambda u^\lambda g_{\alpha\beta} \right) - \zeta \mathcal{P}^{\mu\nu} \partial_\lambda u^\lambda$$

$$\nu^\mu = \sigma \left[\mathcal{P}^{\mu\alpha} \partial_\alpha \left(\frac{\mu}{T} \right) - E^\mu \right]$$

- **Projector** $\mathcal{P}^{\mu\nu} = g^{\mu\nu} + u^\mu u^\nu$
- **Local electric field** $E^\mu = F^{\mu\nu} u_\nu$
- **Transport coefficients: shear, bulk viscosities, conductivity**
- **Entropy production**

$$s^\mu = s u^\mu - \frac{\mu}{T} \nu^\mu$$

$$\partial_\mu s^\mu \geq 0 \quad \Rightarrow \quad \eta \geq 0 \quad \zeta \geq 0 \quad \sigma \geq 0$$

Hydro with anomalies

Conservation laws: effective action with sources $\Gamma[A_\mu, g_{\mu\nu}]$

Gauge Trafo: $\delta A_\mu = \partial_\mu \lambda$ Anomaly

Diffeo Trafo: $\delta g_{\mu\nu} = \nabla_\mu \epsilon_\nu + \nabla_\nu \epsilon_\mu$ $\delta A_\mu = \nabla_\mu (\epsilon^\nu A_\nu) + \epsilon^\nu F_{\mu\nu}$

$$\partial_\mu T^{\mu\nu} = F^{\nu\mu} J_\mu - A^\nu \partial_\mu J^\mu$$

Rewrite with covariant current

$$\begin{aligned} \partial_\mu T^{\mu\nu} &= F^{\nu\mu} J_\mu \\ \partial_\mu J^\mu &= \delta C_{\text{cov}} E^\mu B_\mu \end{aligned}$$

$$E^\mu = F^{\mu\nu} u_\nu, \quad B^\mu = \frac{1}{2} \epsilon^{\mu\nu\rho\lambda} u_\nu F_{\rho\lambda}$$

Hydro with anomalies

$$\partial_\mu T^{\mu\nu} = F^{\nu\mu} J_\mu$$

$$\partial_\mu J^\mu = 8 C_{\text{cov}} E^\mu B_\mu$$

Anomaly breaks parity: additional terms are allowed

$$\delta\mathcal{T}^{\mu\nu} = \sigma_\epsilon^B (u^\mu B^\nu + u^\nu B^\mu) + \sigma_\epsilon^\omega (u^\mu \omega^\nu + u^\nu \omega^\mu)$$

$$\delta\mathcal{V}^\mu = \sigma_B B^\mu + \sigma_\omega \omega^\mu$$

$$\delta\mathcal{S}^\mu = X B^\mu + Y \omega^\mu$$

Vorticity: $\omega^\mu = \frac{1}{2} \epsilon^{\mu\nu\rho\lambda} u_\nu \partial_\rho u_\lambda$

Second law almost fixes them: $\partial_\mu \mathcal{S}^\mu \geq 0$

$$\sigma^B = 24 C_{\text{cov}} \mu$$

$$\sigma_\epsilon^B = 12 C_{\text{cov}} \mu^2 + \gamma T^2$$

$$\sigma^\omega = 24 C_{\text{cov}} \mu^2 + 2\gamma T^2$$

$$\sigma_\epsilon^\omega = 8 C_{\text{cov}} \mu^3 + 2\gamma \mu T^2$$

Hydro with anomalies

- Anomaly+Entropy fix dependence on chemical potentials
- Temperature enters with an undetermined integration constant
- Gravitational anomaly contribution is 4th order in derivatives, naively: no hydro
- **Dissipationless**



Kubo formulas

gravitomagnetic field = metric component

$$ds^2 = -dt^2 + 2A_i^g dt dx^i + d\vec{x}^2$$

$$B_i^g = \epsilon_{ijk} \partial_j A_k^g$$

Relation to vorticity $u^\mu = (1, 0, 0, 0)$, $u_\mu = (-1, A_i^g)$

$$2\vec{\omega} = \vec{B}_i^g$$

Chiral vortical effect is chiral gravitomagnetic effect
(frame dragging, Thirring-Lense effect)

Kubo formulas

Conductivity: $\langle J_i \rangle = \sigma(-i\omega) A_i \implies \sigma = \lim_{\omega \rightarrow 0} \frac{i}{\omega} \langle J_i J_i \rangle$

CME: $\langle J_i \rangle = \sigma_B \epsilon_{ijk} (ip_j) A_j \implies \sigma_B = \lim_{p_z \rightarrow 0} \frac{-i}{p_z} \langle J_x J_y \rangle$

CVE: $\langle J_i \rangle = \frac{1}{2} \sigma_\omega \epsilon_{ijk} (ip_j) A_j^g \implies \sigma_\omega = 2 \lim_{p_z \rightarrow 0} \frac{-i}{p_z} \langle J_x T_{0y} \rangle$

CME in energy current: $\sigma_\epsilon^B = \lim_{p_z \rightarrow 0} \frac{-i}{p_z} \langle T_{0x} J_y \rangle$

$$2\sigma_\epsilon^B = \sigma_\omega$$

Holography

- Hydro has undetermined integration constant
- Weak coupling suggests relation to gravitational anomaly
- Answer with help of holographic model

AdS/CFT

Motto:

“... if the gravitational field didn't exist, one could invent it for the purposes of this paper..”


“Theory of Thermal Transport Coefficients”
Luttinger Phys. Rev. 135, A1505, (1964)

AdS/CFT

“... if string theory didn't exist, one could invent it for the purposes of computing transport coefficients in strongly coupled theories...”

AdS/CFT

$$ds^2 = \frac{r^2}{L^2} (dt^2 + d\vec{x}^2) + \frac{L^2 dr^2}{r^2}$$



AdS/CFT

$$\int_{\Phi|_{\partial}=\Phi_0} D\Phi e^{iS[\Phi]} = e^{iZ[\Phi_0]}$$

$$\frac{\delta^n Z[\Phi_0]}{\delta\Phi_0^1(x_1) \cdots \delta\Phi_0^n(x_n)} = \langle O_1(x_1) \cdots O_n(x_n) \rangle$$

Path integral (string theory) on AdS is hard. In practice resort to semi classical limit:

$$S_{grav}[\Phi_0] = Z[\Phi_0]$$

AdS/CFT

- N=4 SYM best understood example: $\{A_\mu, \Psi_\alpha^a, \phi^I\}$

$$g_{YM}^2 N = \frac{R^4}{\alpha'^2} \quad \frac{1}{N} \propto g_s$$

- *semiclassical gravity limit = large N, large coupling*
- *Interpretation of gravity solutions*

$$\Phi = \Phi_0 (r^{\Delta_-} + \dots) + \langle \mathcal{O} \rangle (r^{\Delta_+} + \dots)$$

- Non-normalizable
- Coupling in dual QFT

- Normalizable
- Vev in dual QFT

AdS/CFT

Dictionary

AdS

five dimensional

r-direction

strongly coupled

gravity

metric

gauge field

scalar field

Field Theory

four dimensional

RG flow

weakly coupled





no gravity

energy momentum tensor

current

scalar operator

String Theory as spherical cow of sQGP



AdS/CFT

- Holographic Model $S = S_{MEH} + S_{CS} + S_{GH} + S_{CSK}$

$$S_{MEH} = \int d^5x \sqrt{-g} \left[\frac{1}{2\kappa^2} (\mathcal{R} + 2\Lambda) - \frac{1}{4} F^2 \right]$$

$$S_{CS} = \int d^5x \sqrt{-g} \epsilon^{MNPQR} A_M \left(\frac{\alpha}{3} F_{NP} F_{QR} + \lambda \mathcal{R}^A{}_{BNP} \mathcal{R}^B{}_{AQR} \right)$$

$$S_{GH} = \int_{\partial} d^4x \sqrt{-h} \frac{1}{\kappa^2} K$$

$$S_{CSK} = \int_{\partial} d^4x \sqrt{-h} \frac{4}{\kappa^2} \lambda n_M \epsilon^{MNPQR} A_N K_{PL} D_Q K_R^L$$

AdS/CFT

- Anomaly: $\delta S = \int_{\partial} d^4x \sqrt{-h} \epsilon^{m n k l} \left(\frac{\alpha}{3} F_{m n} F_{k l} + \lambda \mathcal{R}^a{}_{b m n} \mathcal{R}^b{}_{a k l} \right)$

- Background:

$$ds^2 = \frac{r^2}{L^2} \left(-f(r) dt^2 + d\vec{x}^2 \right) + \frac{L^2}{r^2} dr^2 \quad f = 1 - \frac{M}{r^4} + \frac{Q}{r^6}$$

$$A = \left(A_0 - \frac{\mu r_H^2}{r^2} \right) dt$$

allow for general boundary value !

- Definition of Currents:

$$J^\mu = \sqrt{-h} \frac{1}{2\kappa^2} F^{\mu r} + \frac{4\alpha}{3} \epsilon^{m n k l} A_n F_{k l}$$

consistent current

covariant current

Bardeen-Zumino polynomial

AdS/CFT

- Chiral Gravitomagnetic Effect via AdS boundary conditions

$$\delta h^t{}_x|_{r=\infty} = B_g y \quad , \quad \delta A_x|_{r=\infty} = -\mu u y B_g$$

- Response parallel to gravitomagnetic field

$$[f a'_z - \mu h^t{}_z]' = -\alpha 4\mu u B_g + \lambda 4f' (2uf'' + f') B_g$$

- Induced covariant current

$$\vec{J}_{\text{cov}} = (4\alpha\mu + 32\lambda T^2) \vec{B}_g$$

- Normalize to anomaly of 1 chiral fermion

$$\alpha = \frac{1}{96\pi^2} \quad , \quad \lambda = \frac{1}{768\pi^2}$$

- Matches precisely the weak coupling result!!


4 derivatives !

Ads/CFT



Remarks

- Dynamical Gauge fields:
radiative corrections due to rescattering
- Non-renormalization holds for 't Hooft anomalies
(external gauge fields, anomalies with only global symmetries)
- Holographic model: Stückelberg axion
- Anomalous dimension of axial current = mass term for gauge field in AdS



$$\int d^5x \sqrt{-g} \left(-\frac{1}{4} F^2 + m^2 (A_\mu + \partial_\mu \theta) + \frac{\alpha}{3} (A + d\theta) \wedge F \wedge F \right)$$

$$\partial_\mu J_5^\mu = \frac{\delta Z}{\delta \theta}$$

- Axial charge decay but basic transport phenomena persist

Theory Summary

$$\vec{J} = \frac{\mu_5 - A_0^5}{2\pi^2} \vec{B} \quad \text{Chiral Magnetic Effect (CME)}$$


$$\vec{J}_5 = \frac{\mu}{2\pi^2} \vec{B} \quad \text{Chiral Separation Effect (CSE)}$$

$$\vec{J} = \frac{2\mu\mu_5}{\pi^2} \vec{\Omega} \quad \text{Chiral Vortical Effects (CVE)}$$

$$\vec{J}_5 = \left(\frac{\mu^2 + \mu_5^2}{\pi^2} + \frac{T^2}{12} \right) \vec{\Omega}$$

Application in QGP

Induced Quadrupole moment in HICs



2 steps


$$\vec{J}_5 = \frac{\mu}{2\pi^2} \vec{B}$$

$$\vec{J} = \frac{\mu_5}{2\pi^2} \vec{B}$$

propagating wave of axial-
electric charge conversion:
Chiral Magnetic Wave

Application in QGP

Induced Quadrupole moment in HICs



[Star Collaboration (RHIC)]

2 steps

$$\vec{J}_5 = \frac{\mu}{2\pi^2} \vec{B}$$

$$\vec{J} = \frac{\mu_5}{2\pi^2} \vec{B}$$

propagating wave of axial-
electric charge conversion:
Chiral Magnetic Wave

Summary

- Hydrodynamics with anomalies
- Local form of 2nd law almost fixes CME, CVE
- Derivative mismatch for grav. anomaly in CVE avoided in Holography (additional direction)
- (Non)-Renormalization
- Applications: CMV

Outlook

- Weyl Semi-metals and CME and NMR
- Edge physics (Fermi arcs) from CVE and CME
- Anomalous Hall Effect (AHE)
- Holographic model of WSM