

Quasi-distribution amplitudes for pion and kaon via the nonlocal chiral-quark model (NLChQM)

Nam, Seung-il

Department of Physics and
Institute for Radiation Science & Technology (IRST),
Pukyong National University (PKNU),
Busan, Republic of Korea




Contents based on **SiN, Modern Physics Letters A32, 1750218 (2017)**

Theory


Quark distribution amplitude (DA) for PS mesons

Pion-photon transition form factor




A.V.Radyushkin, PRD80, 094009 (2009)

Hard exclusive pion electroproduction




K.Park et al. [CLAS Collaboration], PLB780, 340 (2018)

Heavy-meson weak decay



Structural (nonpert.) information of DA as func. of x

Theory



PS-meson DA defined in light-front (LF) formalism

$$\langle 0 | \bar{q}_f(\tau \hat{n}) \gamma_\mu \gamma_5 q_g(-\tau \hat{n}) | \mathcal{M}(P) \rangle = i\sqrt{2} F_{\mathcal{M}} P_\mu \int_0^1 du e^{i(2u-1)P \cdot \tau \hat{n}} \phi_{\mathcal{M}}(u),$$

Well defined Fock-states for meson but not covariant

LF formalism can not applied for lattice QCD (LQCD)

Instead, LQCD computes *moments* of DA

X.Ji, A.V.Radyushkin have developed quasi-DA (QDA) in terms of Large-momentum effective theo. (LaMET)

Theory (LaMET) A. V. Radyushkin, PRD93, 056002 (2016).

Defining virtuality distribution amplitude (VDA)

$$\langle 0 | \bar{q}(0) \gamma_5 \gamma_+ q(z_-) | \mathcal{M}(p) \rangle = \frac{p_+}{\sqrt{2} F_{\mathcal{M}}} \int_0^\infty d\sigma \int_0^1 dx \Phi_{\mathcal{M}}(x, \sigma) e^{-ixp_+ z_-}.$$

Relation between DA and VDA

$$\phi_{\mathcal{M}}(x) = \int_0^\infty d\sigma \Phi_{\mathcal{M}}(x, \sigma), \quad \int_0^1 dx \phi_{\mathcal{M}}(x) = 1$$

Fourier transform (FT) of matrix element: TMDA

$$\langle 0 | \bar{q}(0) \gamma_5 \gamma_+ q(z_-) | \mathcal{M}(p) \rangle = \frac{p_+}{\sqrt{2} F_{\mathcal{M}}} \int_0^\infty d^2 k_\perp \int_0^1 dx \Psi_{\mathcal{M}}(x, k_\perp^2) e^{-ixp_+ z_-}.$$

Theory (LaMET)

TMDA in terms of VDA

$$\Psi_{\mathcal{M}}(x, k_{\perp}^2) = \frac{i}{\pi} \int_0^{\infty} \frac{d\sigma}{\sigma} \Phi_{\mathcal{M}}(x, \sigma) e^{-ik_{\perp}^2/\sigma}$$

TMDA integrated over k_{\perp} gives DA

$$\phi_{\mathcal{M}}(x) = \int_0^{\infty} dk_{\perp}^2 \Psi_{\mathcal{M}}(x, k_{\perp}^2) = 2\pi \int_0^{\infty} k_{\perp} dk_{\perp} \Psi_{\mathcal{M}}(x, k_{\perp}^2)$$

DA

$$\text{VDA} \Leftrightarrow \text{TMDA} \Leftrightarrow \text{DA}$$

Theory (LaMET)

Now, matrix element at equal time

$$z = (0, 0, 0, z_3)$$

$$\langle 0 | \bar{q}(0) \gamma_5 \gamma_0 q(z_3) | \mathcal{M}(p) \rangle = \frac{p_0}{\sqrt{2} F_{\mathcal{M}}} \int_0^{\infty} d\sigma \int_0^1 dx \Phi_{\mathcal{M}}(x, \sigma) e^{-ixp_3 z_3 + i\sigma z_3^2/4}.$$

VDA

Similarly, FT of equal-time matrix element: QDA $[-\infty, \infty]$

$$\langle 0 | \bar{q}(0) \gamma_5 \gamma_0 q(z_3) | \mathcal{M}(p) \rangle = \frac{p_0}{\sqrt{2} F_{\mathcal{M}}} \int_{-\infty}^{\infty} dy Q_{\mathcal{M}}(y, p_3) e^{-iy p_3 z_3}.$$

QDA

QDA in terms of VDA: Constrained x , while not for y

$$Q_{\mathcal{M}}(y, p_3) = \int_0^1 dx \int_0^{\infty} d\sigma \sqrt{\frac{ip_3^2}{\pi\sigma}} e^{-i(x-y)^2 p_3^2/\sigma} \Phi_{\mathcal{M}}(x, \sigma),$$

Theory (LaMET)

By equating them, TMDA and VDA related as

$$p_3 \int_{-\infty}^{\infty} dk_1 \Psi_{\mathcal{M}}(x, k_1^2 + (x-y)^2 p_3^2) = \int_0^{\infty} d\sigma \sqrt{\frac{ip_3^2}{\pi\sigma}} e^{-i(x-y)^2 p_3^2/\sigma} \Phi_{\mathcal{M}}(x, \sigma).$$

Thus, QDA given in terms of TMDA A. V. Radyushkin, PRD93, 056002 (2016).

$$Q_{\mathcal{M}}(y, p_3) = p_3 \int_{-\infty}^{\infty} dk_1 \int_0^1 dx \Psi_{\mathcal{M}}(x, k_1^2 + (x-y)^2 p_3^2).$$

A useful limit $p_3 \rightarrow \infty$

$$\lim_{p_3 \rightarrow \infty} \left[\sqrt{\frac{ip_3^2}{\pi\sigma}} e^{-i(x-y)^2 p_3^2/\sigma} \right] = \delta(x-y)$$

Theory (LaMET)

Due to the limit, DA-like func. (covariant) relates to VDA

$$\begin{aligned} \lim_{p_3 \rightarrow \infty} Q_{\mathcal{M}}(y, p_3) &\equiv \overset{\sim\text{DA}}{\varphi_{\mathcal{M}}(y)} = \int_0^1 dx \int_0^\infty d\sigma \delta(x - y) \Phi_{\mathcal{M}}(x, \sigma) \\ &= \int_0^\infty d\sigma \Phi_{\mathcal{M}}(y, \sigma). \end{aligned}$$

DA-like func. and DA satisfy similar normalizations

$$\int_{-\infty}^{\infty} dy \varphi_{\mathcal{M}}(y) = \int_0^1 dx \phi_{\mathcal{M}}(x) = 1$$

Theory (LaMET with a model for LFWF)

Introducing LFWF for DA, previous equation becomes

$$\lim_{p_3 \rightarrow \infty} p_3 \int_{-\infty}^{\infty} dy \int_{-\infty}^{\infty} dk_1 \int_{-\infty}^{\infty} dk_2 \Psi_{\mathcal{M}}(x, k_{\perp}^2) \delta(k_2 - (x - y)p_3)$$

TMDA

$$= \int_0^{\infty} d^2 k_{\perp} \psi_{\mathcal{M}}(x, k_{\perp}^2)$$

LFWF

After performing integration, arriving at

$$\lim_{p_3 \rightarrow \infty} \Psi_{\mathcal{M}}(x, k_{\perp}^2) \Big|_{x=y+\frac{k_2}{p_3}} = \psi_{\mathcal{M}}(x, k_{\perp}^2).$$

If k_2 is small (nonpert.), $\lim_{p_3 \rightarrow \infty} k_2/p_3 = 0$


Theory (LaMET with a model for LFWF)

As far as we are interested in NP region, we have

$$\Psi_{\mathcal{M}}^{\text{NP}}(y, k_{\perp}^2) = \psi_{\mathcal{M}}^{\text{NP}}(y, k_{\perp}^2) \quad \text{for } y = x = [0, 1]$$


Saying, LFWF \sim TMDA in NP region

How can we test this relation?


$$F_{\gamma\gamma^*\pi^0}(Q^2) = \frac{2F_{\pi}}{3} \int_0^1 \frac{dx}{xQ^2} \int_0^{xQ^2} \frac{dk_{\perp}^2}{xQ^2} \int_0^{k_{\perp}} d^2p_{\perp} \Psi(x, p_{\perp}^2)$$


Replacing TMDA with LFWF, data reproduced?


Theory (general)



Theory (LaMET)



Theory (LaMET with NLChQM)



Model: Non-local chiral-quark model (NLChQM)

Based on liquid-instanton model (LIM)

D. Diakonov, Prog. Part. Nucl. Phys. 51, 173 (2003).

Nonlocal inter-quark interactions via instanton

$$\mathcal{S}_{\text{eff}}[m_q, \mathcal{M}] = -\text{Sp} \ln \left[i\rlap{\not{D}} + im_q + i\sqrt{M_q(\partial^2)}U^{\gamma_5}(\mathcal{M})\sqrt{M_q(\partial^2)} \right]$$

Effective model describing SCSB at $\Lambda_{\text{NLChQM}} \sim 1.0 \text{ GeV}$

Performing Wick rotation from Euclidean to Minkowski

Then, Minkowski to LF frame with light-like vector n

Model: Non-local chiral-quark model (NLChQM)

PS-meson DA within NLChQM reads

$$\phi_{\mathcal{M}}^{\text{NP}}(x) = -\frac{2iN_c}{F_{q\bar{q}'}^2} \int \frac{d^4k}{(2\pi)^4} \sqrt{M_q(k)M_{q'}(k-p)} \delta[\bar{x}p \cdot n - k \cdot n] \\ \times \frac{[M_{q'}k - M_q(k-p)] \cdot n}{[k^2 - M_q^2][(k-p)^2 - M_{q'}^2]}.$$

SiN et al., Phys. Rev. D 74, 014019 (2006)

And, LFWF from NLChQM reads SiN, MPLA32, 1750218 (2017)

$$\psi_{\mathcal{M}}^{\text{NP}}(x, k_{\perp}^2)$$

$$\bar{x} = 1 - x$$


$$= \frac{\bar{x}N_c\Lambda^4 \sqrt{M_q M_{q'}} [xM_q + \bar{x}M_{q'}]}{4\pi^3 F_{\mathcal{M}}^2 [M_q^2 - \Lambda^2] [k_{\perp}^2 + \Lambda^2 - x\bar{x}M_{\mathcal{M}}^2] [k_{\perp}^2 + \bar{x}M_{q'}^2 + x\Lambda^2 - x\bar{x}M_{\mathcal{M}}^2]}$$

$$+ \frac{\bar{x}N_c\Lambda^4 \sqrt{M_q M_{q'}} [xM_q + \bar{x}M_{q'}]}{4\pi^3 F_{\mathcal{M}}^2 [\Lambda^2 - M_q^2] [k_{\perp}^2 + xM_q^2 + \bar{x}\Lambda^2 - x\bar{x}M_{\mathcal{M}}^2] [k_{\perp}^2 + xM_q^2 + \bar{x}M_{q'}^2 - x\bar{x}M_{\mathcal{M}}^2]},$$

Model: Non-local chiral-quark model (NLChQM)

An **analytic** expression for QDA beyond chiral limit within NLChQM [SiN, MPLA32, 1750218 \(2017\)](#)

$$Q_{\mathcal{M}}^{\text{NP}}(y, p_3) = \frac{N_c M_0^2 \Lambda^4}{8\pi^2 F_\pi^2 \eta^4} \times \left\{ \ln \left[\frac{[\eta^2 + 2p_3 f_-(\bar{y}, \Lambda)]^2 [\eta^2 - 2p_3 f_-(\bar{y}, M_0)]^2 f_+(\bar{y}, \Lambda) f_+(\bar{y}, M_0) f_+(y, \Lambda) f_+(y, M_0)}{[\eta^2 + 2p_3 f_+(y, \Lambda)]^2 [\eta^2 - 2p_3 f_+(y, M_0)]^2 f_-(\bar{y}, \Lambda) f_-(\bar{y}, M_0) f_-(y, \Lambda) f_-(y, M_0)} \right] \right. \\ \left. - \frac{\Delta M}{M_0} \left[(2y - 3) \ln \left[\frac{f_-(\bar{y}, \Lambda) f_-(\bar{y}, M_0) [\eta^2 + 2p_3 f_+(y, \Lambda)] [\eta^2 - 2p_3 f_+(y, M_0)]}{f_+(y, \Lambda) f_+(y, M_0) [\eta^2 + 2p_3 f_-(\bar{y}, \Lambda)] [\eta^2 - 2p_3 f_-(\bar{y}, M_0)]} \right] \right] \right. \\ \left. + \frac{\eta^2}{p_3^2} \ln \left[\frac{[\eta^2 + 2p_3 f_+(y, \Lambda)] [\eta^2 + 2p_3 f_-(\bar{y}, \Lambda)]}{[\eta^2 - 2p_3 f_+(y, M_0)] [\eta^2 - 2p_3 f_-(\bar{y}, M_0)]} \right] \right\} + \mathcal{O}(\Delta M^2),$$



$$\Delta M = |M_q - M_{q'}| = |m_q - m_{q'}|$$

$$f_{\pm}(y, M_0) = xp_3 \pm \sqrt{M_0^2 + y^2 p_3^2},$$

$$\eta^2 = \Lambda^2 - M_0^2$$


Numerical results

Table 1. Model parameters for the present calculations. With these values, the pion and kaon DAs satisfy the normalization condition, i.e. $\int dx \phi_{\pi,K}(x) = 1$.

	$M_0 = 350 \text{ MeV}$	$m_{u,d} = 5 \text{ MeV}$	$m_s = 135 \text{ MeV}$	
Pion at CL	$F_\pi = 93 \text{ MeV}$	$M_q = M_0$	$M_{q'} = M_0$	$M_\pi = 0 \text{ MeV}$
Pion	$F_\pi = 93 \text{ MeV}$	$M_q = (m_{u,d} + M_0)$ $= 355 \text{ MeV}$	$M_{q'} = (m_{u,d} + M_0)$ $= 355 \text{ MeV}$	$M_\pi = 140 \text{ MeV}$
Kaon	$F_K = 113 \text{ MeV}$	$M_q = (m_{u,d} + M_0)$ $= 355 \text{ MeV}$	$M_{q'} = (m_s + M_0)$ $= 485 \text{ MeV}$	$M_K = 495 \text{ MeV}$


$\Lambda = 1.02 \text{ GeV}$
 $\Lambda = 1.01 \text{ GeV}$
 $\Lambda = 1.05 \text{ GeV}$

TMDA ~ LFWF as a func. of (x, k_T)



SiN, MPLA32, 1750218 (2017)


DA as a func. of x



Numerical results

$$F_{\gamma\gamma^*\pi^0}(Q^2) = \frac{2F_\pi}{3} \int_0^1 \frac{dx}{xQ^2} \int_0^{xQ^2} \frac{dk_\perp^2}{xQ^2} \int_0^{k_\perp} d^2p_\perp \Psi(x, p_\perp^2).$$

Photon-pion transition FF with TMDA ~ LFWF



Numerical results

Non-zero at $Q^2=0$, via Adler-Bell-Jackiw axial anomaly for real photons: $F_{\gamma\gamma\pi^0}(0) = (4\pi^2 F_\pi)^{-1} \approx 0.272 \text{ GeV}^{-1}$

From NLChQM gives $\lim_{Q^2 \rightarrow 0} F_{\gamma\gamma^*\pi^0}^{\text{NLChQM}}(Q^2) = 0.191 \text{ GeV}^{-1}$
SiN, MPLA32, 1750218 (2017)


The ratio becomes $F_{\gamma\gamma^*\pi^0}^{\text{NLChQM}}(0)/F_{\gamma\gamma\pi^0}(0) = 0.702$

Slightly larger than 0.5 G. P. Lepage and S. J. Brodsky, PRD22, 2157 (1980)
A. V. Radyushkin, PRD93, 056002 (2016).

E. Ruiz Arriola and W. Broniowski, PRD74, 034008 (2006)
A. G. Oganesian et al., PRD93, 054040 (2016).

Numerical results


Pion and kaon QDA for p_3 [GeV]



Slightly tilted curves for kaon, due to $m_s > m_u, m_d$

Numerical results


Pion QDA for p_3 [GeV] and y



Numerical results

J. Gronberg et al. (CLEO Collaboration), Phys. Rev. D 57, 33 (1998).
A. Schmedding and O. I. Yakovlev, Phys. Rev. D 62, 116002 (2000).

Gegenbauer coefficients from pion QDA for $p_3 \rightarrow \infty$



Numerical results

Moments from DA and QDA $\xi \equiv (y - \bar{y}) = 2x - 1$

$$\langle \xi^n \rangle_{\mathcal{M}}^{\text{DA}} = \int_0^1 (2x - 1)^n \phi_{\mathcal{M}}(x) dx, \quad \langle \xi^n \rangle_{\mathcal{M}}^{\text{QDA}} = \lim_{p_3 \rightarrow \infty} \int_{-\infty}^{\infty} (2y - 1)^n Q_{\mathcal{M}}(y, p_3) dy$$

Table 2. The moments from the pion and kaon DA and QDA for $\xi = 2x - 1$ and $2y - 1$, respectively.

	$n = 1$	$n = 2$	$n = 3$	$n = 4$		$n = 1$	$n = 2$	$n = 3$	$n = 4$
$\langle \xi^n \rangle_{\pi}^{\text{DA}}$	–	0.2210	–	0.1002	$\langle \xi^n \rangle_K^{\text{DA}}$	0.0277	0.2043	0.0122	0.0887
$\langle \xi^n \rangle_{\pi}^{\text{QDA}}$ at $p_3 = 10\Lambda$	–	0.2287	–	0.1159	$\langle \xi^n \rangle_K^{\text{QDA}}$ at $p_3 = 10\Lambda$	0.0277	0.2118	0.0120	0.1034
$\langle \xi^n \rangle_{\pi}^{\text{QDA}}$ at $p_3 = 20\Lambda$	–	0.2229	–	0.1030	$\langle \xi^n \rangle_K^{\text{QDA}}$ at $p_3 = 20\Lambda$	0.0277	0.2062	0.0121	0.0913
$\langle \xi^n \rangle_{\pi}^{\text{QDA}}$ at $p_3 = 30\Lambda$	–	0.2218	–	0.1013	$\langle \xi^n \rangle_K^{\text{QDA}}$ at $p_3 = 30\Lambda$	0.0277	0.2052	0.0122	0.0898


$$\langle \xi^2 \rangle_{\pi}^{\overline{\text{MS}}} = \int_0^1 du (2u - 1)^2 \phi_{\pi}(u, \mu) = 0.2361(41)(39)(?)$$

V.Braun et al. (RQCD Collaboration), PRD 92, 014504 (2015)

For $p_3 > 30$ GeV, differences reduced to $\sim 10\%$

Numerical results

Behavior of moments for p_3



Higher moments depend much on p_3

Summary and perspectives

Verified that TMDA \sim LFWF at NP region

From LFWF from NLCHQM, analytic form for QDA derived for nonzero current-quark mass

Obtained QDA for pion and kaon successfully describe DAs for $p_3 \rightarrow \infty$, showing reproduction of exp. data

Higher moments are sensitive for p_3

Fragmentation func. (FragF) \leftrightarrow PDF via Drell-Levy-Yan
Then, QPDF \leftrightarrow QFragF ?!?! **In progress!!**

Night view of Busan



Thank you for your attention!

This talk supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government (MSIT) (No. 2018R1A5A1025563)