

Skyrmions with low binding energies

Martin Speight (Leeds)

joint with

Derek Harland, Ben Maybee (Leeds)

Mike Gillard (Loughborough)

Elliot Kirk (Durham)

1st APCTP-TRIUMF Joint Workshop on Understanding Nuclei
from Different Theoretical Approaches September 14-19, 2018
APCTP, Pohang, Korea

The (extended) Skyrme model

$$U : \mathbb{R}^3 \rightarrow SU(2), \quad U(\infty) = \mathbb{I}_2, \quad L_i = U^\dagger \partial_i U$$

$$B(U) = \frac{1}{2\pi^2} \int_{\mathbb{R}^3} \underbrace{\frac{\varepsilon_{ijk}}{12} \text{tr}(L_i L_j L_k)}_{\mathcal{B}} \in \mathbb{Z}$$

The (extended) Skyrme model

$$U : \mathbb{R}^3 \rightarrow SU(2), \quad U(\infty) = \mathbb{I}_2, \quad L_i = U^\dagger \partial_i U$$

$$B(U) = \frac{1}{2\pi^2} \int_{\mathbb{R}^3} \underbrace{\frac{\varepsilon_{ijk}}{12} \text{tr}(L_i L_j L_k)}_{\mathcal{B}} \in \mathbb{Z}$$

$$E(U) = \underbrace{-\frac{1}{4} \int_{\mathbb{R}^3} \text{tr}(L_i L_i)}_{E_2} - \underbrace{\frac{1}{8} \int_{\mathbb{R}^3} \text{tr}([L_i, L_j][L_i, L_j])}_{E_4}$$

The (extended) Skyrme model

$$U : \mathbb{R}^3 \rightarrow SU(2), \quad U(\infty) = \mathbb{I}_2, \quad L_i = U^\dagger \partial_i U$$

$$B(U) = \frac{1}{2\pi^2} \int_{\mathbb{R}^3} \underbrace{\frac{\varepsilon_{ijk}}{12} \text{tr}(L_i L_j L_k)}_{\mathcal{B}} \in \mathbb{Z}$$

$$E(U) = \underbrace{-\frac{1}{4} \int_{\mathbb{R}^3} \text{tr}(L_i L_i)}_{E_2} - \underbrace{\frac{1}{8} \int_{\mathbb{R}^3} \text{tr}([L_i, L_j][L_i, L_j])}_{E_4} + \underbrace{\int_{\mathbb{R}^3} V(U)}_{E_0} + \underbrace{\frac{1}{2} \int_{\mathbb{R}^3} \mathcal{B}^2}_{E_6}$$

E minimizer of charge B : classical model of nucleon number B nucleus

The binding energy problem ($E = E_2 + E_4$)

$$\text{Classical binding energy} = \frac{BE(U_1) - E(U_B)}{E(U_1)}$$

| B | Element | B.E. (Skyrme) | B.E. (experiment) |
|-----|---------|---------------|-------------------|
| 4 | He | 0.3639 | 0.0301 |
| 7 | Li | 0.7811 | 0.0414 |
| 9 | Be | 1.0123 | 0.0615 |
| 11 | B | 1.2792 | 0.0807 |
| 12 | C | 1.4277 | 0.0981 |
| 14 | N | 1.6815 | 0.1114 |
| 16 | O | 1.9646 | 0.1359 |
| 19 | F | 2.3684 | 0.1570 |
| 20 | Ne | 2.5045 | 0.1710 |

Naive quantization makes the problem **worse**

A solution?

- Faddeev showed that $E = E_2 + E_4$ has a topological lower bound:

$$E \geq \text{const} \times B$$

Unfortunately (?) it's never attained

A solution?

- Faddeev showed that $E = E_2 + E_4$ has a topological lower bound:

$$E \geq \text{const} \times B$$

Unfortunately (?) it's never attained

- What if we find a Skyrme energy with a bound like this which is attained for each B ?
- Then $E(U_B) \equiv B \times E(U_1)$ so $B.E. \equiv 0!$
- Call such a model "BPS"

A solution?

- Faddeev showed that $E = E_2 + E_4$ has a topological lower bound:

$$E \geq \text{const} \times B$$

Unfortunately (?) it's never attained

- What if we find a Skyrme energy with a bound like this which is attained for each B ?
- Then $E(U_B) \equiv B \times E(U_1)$ so $B.E. \equiv 0!$
- Call such a model “BPS”
- Now perturb the BPS model by adding $\varepsilon(E_2 + E_4)$, ε small (for example)
- Hopefully get a “near BPS” Skyrme model with small positive BEs

A solution?

- Faddeev showed that $E = E_2 + E_4$ has a topological lower bound:

$$E \geq \text{const} \times B$$

Unfortunately (?) it's never attained

- What if we find a Skyrme energy with a bound like this which is attained for each B ?
- Then $E(U_B) \equiv B \times E(U_1)$ so $B.E. \equiv 0!$
- Call such a model “BPS”
- Now perturb the BPS model by adding $\varepsilon(E_2 + E_4)$, ε small (for example)
- Hopefully get a “near BPS” Skyrme model with small positive BEs
- Two implementations of this idea
 - perturbed $E_6 + E_0$ model (Adam, Sanchez-Guillen, Wereszczynski)
 - perturbed $E_4 + E_0$ model (Harland)

The ASW model ($E_6 + E_0$)

- $U : M \rightarrow N$, $\Omega =$ volume form on N
($M = \mathbb{R}^3$, $N = SU(2) = S^3$)
- Potential $V = \frac{1}{2} W^2$ where $W : N \rightarrow \mathbb{R}$ has $W(\mathbb{I}_2) = 0$

$$E(U) = \frac{1}{2} \int_M (|U^* \Omega|^2 + W(U)^2)$$

The ASW model ($E_6 + E_0$)

- $U : M \rightarrow N$, $\Omega = \text{volume form on } N$
($M = \mathbb{R}^3$, $N = SU(2) = S^3$)
- Potential $V = \frac{1}{2} W^2$ where $W : N \rightarrow \mathbb{R}$ has $W(\mathbb{I}_2) = 0$

$$E(U) = \frac{1}{2} \int_M (|U^* \Omega|^2 + W(U)^2)$$

- Energy bound

$$\begin{aligned} 0 &\leq \frac{1}{2} \int_M (*U^* \Omega - W(U))^2 = E - \int_M U^*(W\Omega) \\ &= E - \underbrace{\langle W \rangle \text{Vol}(N)}_C B \end{aligned}$$

- So $E \geq CB$ with equality iff $U^* \Omega = *W(U)$

$$U : M \rightarrow N, \quad U^* \Omega = *W(U)$$

$$U : M \rightarrow N, \quad U^* \Omega = *W(U)$$

$$U : M' \rightarrow N', \quad U^* \left(\frac{\Omega}{W} \right) = *1 = \text{volume form on } M$$

$N' = N \setminus W^{-1}(0)$ = punctured target space,

$M' = U^{-1}(N')$ = “support” of U

$$U : M \rightarrow N, \quad U^* \Omega = *W(U)$$

$$U : M' \rightarrow N', \quad U^* \left(\frac{\Omega}{W} \right) = *1 = \text{volume form on } M$$

$N' = N \setminus W^{-1}(0)$ = punctured target space,

$M' = U^{-1}(N')$ = "support" of U

- BPS skyrmions are **volume preserving** maps

$$M' \rightarrow (N', \Omega/W)$$

$$U : M \rightarrow N, \quad U^* \Omega = *W(U)$$

$$U : M' \rightarrow N', \quad U^* \left(\frac{\Omega}{W} \right) = *1 = \text{volume form on } M$$

$N' = N \setminus W^{-1}(0)$ = punctured target space,

$M' = U^{-1}(N')$ = "support" of U

- BPS skyrmions are **volume preserving** maps
 $M' \rightarrow (N', \Omega/W)$
- Come in ∞ -dim families, $U \circ \psi$, where $\psi : M \rightarrow M$ is any volume preserving map. Cf Liquid drop model.

$$U : M \rightarrow N, \quad U^* \Omega = *W(U)$$

$$U : M' \rightarrow N', \quad U^* \left(\frac{\Omega}{W} \right) = *1 = \text{volume form on } M$$

$N' = N \setminus W^{-1}(0)$ = punctured target space,

$M' = U^{-1}(N')$ = "support" of U

- BPS skyrmions are **volume preserving** maps
 $M' \rightarrow (N', \Omega/W)$
- Come in ∞ -dim families, $U \circ \psi$, where $\psi : M \rightarrow M$ is any volume preserving map. Cf Liquid drop model.
- Compactons? Depends on W . Support of U_1 has

$$\text{Vol} = \int_{N'} \frac{\Omega}{W}$$

$$U^* \Omega = *W(U)$$

- $B = 1$ Hedgehog (assume $W = W(\text{tr } U)$, preserves chiral symmetry)

$$U_H(r\mathbf{n}) = \cos f(r) + i \sin f(r) \mathbf{n} \cdot \boldsymbol{\tau} \quad f(0) = \pi, f(\infty) = 0$$

1st order ODE for f .

$$U^* \Omega = *W(U)$$

- $B = 1$ Hedgehog (assume $W = W(\text{tr } U)$, preserves chiral symmetry)

$$U_H(r\mathbf{n}) = \cos f(r) + i \sin f(r) \mathbf{n} \cdot \boldsymbol{\tau} \quad f(0) = \pi, f(\infty) = 0$$

1st order ODE for f .

- Charge B solutions (ASW, Bonenfant, Marleau)

$$\psi_B : \mathbb{R}^3 \setminus \mathbb{R}_z \rightarrow \mathbb{R}^3 \setminus \mathbb{R}_z, \quad \psi_B(r, \theta, \varphi) = (B^{-1/3}r, \theta, B\varphi)$$

$U_B = U_H \circ \psi_B$. Conical singularity along \mathbb{R}_z

Perturbation

- $E_6 + E_0$ is irredeemably sick (e.g. EL eqn isn't an evolution eqn)
- Need $E = E_6 + E_0 + \varepsilon E_2$

Perturbation

- $E_6 + E_0$ is irredeemably sick (e.g. EL eqn isn't an evolution eqn)
- Need $E = E_6 + E_0 + \varepsilon E_2$
- Pion mass? Linearize about vacuum $U = \mathbb{I}_2 + i\pi \cdot \tau + \dots$

$$V(U) = \frac{1}{2}m^2|\pi|^2 + \dots$$

$$E_{lin} = \int_M \left(\frac{\varepsilon}{2} \partial_i \pi \cdot \partial_i \pi + \frac{m^2}{2} |\pi|^2 \right)$$

Klein-Gordon triplet of mass $m/\sqrt{\varepsilon}$

Perturbation

- $E_6 + E_0$ is irredeemably sick (e.g. EL eqn isn't an evolution eqn)
- Need $E = E_6 + E_0 + \varepsilon E_2$
- Pion mass? Linearize about vacuum $U = \mathbb{I}_2 + i\pi \cdot \tau + \dots$

$$V(U) = \frac{1}{2}m^2|\pi|^2 + \dots$$

$$E_{lin} = \int_M \left(\frac{\varepsilon}{2} \partial_i \pi \cdot \partial_i \pi + \frac{m^2}{2} |\pi|^2 \right)$$

Klein-Gordon triplet of mass $m/\sqrt{\varepsilon}$

- Better choose V with $m = 0$, else pions are heavier than nucleons!
- $V_\pi = \text{tr}(\mathbb{I}_2 - U)$ no good
- $W = [\text{tr}(\mathbb{I}_2 - U)]/2$ is OK. Compacton at $\varepsilon = 0$

Numerical results $E = E_6 + E_0 + \varepsilon E_2$

- Numerics (Harland, Gillard, JMS):
 - Start at $\varepsilon = 1$, minimize using conjugate gradient method.
 - Reduce ε , repeat
 - Check integrality of B and Derrick scaling identities

$$\begin{aligned} E(U(\lambda \mathbf{x})) &= \lambda^3 E_6 + \varepsilon \lambda^{-1} E_2 + \lambda^{-3} E_0 \\ \Rightarrow 0 &= 3E_6 - \varepsilon E_2 - 3E_0 \end{aligned}$$

Numerical results $E = E_6 + E_0 + \varepsilon E_2$

- Numerics (Harland, Gillard, JMS):
 - Start at $\varepsilon = 1$, minimize using conjugate gradient method.
 - Reduce ε , repeat
 - Check integrality of B and Derrick scaling identities

$$\begin{aligned} E(U(\lambda \mathbf{x})) &= \lambda^3 E_6 + \varepsilon \lambda^{-1} E_2 + \lambda^{-3} E_0 \\ \Rightarrow 0 &= 3E_6 - \varepsilon E_2 - 3E_0 \end{aligned}$$

- Numerics become unstable at $\varepsilon \approx 0.2$.
- B.E.s decrease with ε , but still too large

Numerical results $E = E_6 + E_0 + \varepsilon E_2$

- Numerics (Harland, Gillard, JMS):
 - Start at $\varepsilon = 1$, minimize using conjugate gradient method.
 - Reduce ε , repeat
 - Check integrality of B and Derrick scaling identities

$$\begin{aligned} E(U(\lambda \mathbf{x})) &= \lambda^3 E_6 + \varepsilon \lambda^{-1} E_2 + \lambda^{-3} E_0 \\ \Rightarrow 0 &= 3E_6 - \varepsilon E_2 - 3E_0 \end{aligned}$$


- Numerics become unstable at $\varepsilon \approx 0.2$.
- B.E.s decrease with ε , but still too large
- $B = 1, 2$ have axial symmetry: can push ε much further

$$\frac{2E(1) - E(2)}{2E(1)} \approx 0.01$$

requires $\varepsilon = 0.014$, way too small for 3D numerics

Numerical results $E = E_6 + E_0 + \varepsilon E_2$


$$\mathcal{B} = 0.5\mathcal{B}_{max}$$



Left $\varepsilon = 1$, right $\varepsilon = 0.2$

Numerical results $E = E_6 + E_0 + \varepsilon E_2$

Covergence to BPS skyrmions? $B = 4$



Restricted harmonicity

$$E(U) = E_6(U) + E_0(U) + \varepsilon E_2(U)$$

- $E(U)$ should be stationary for all smooth variations U_t of U
- Choose $U_t = U \circ \psi_t$, ψ_t a curve through Id in $S\text{Diff}(M)$

$$E(U) = E_6(U) + E_0(U) + \varepsilon E_2(U)$$

- $E(U)$ should be stationary for all smooth variations U_t of U
- Choose $U_t = U \circ \psi_t$, ψ_t a curve through Id in $SDiff(M)$
- E_6 and E_0 are $SDiff$ invariant! So E_2 must be stationary for these variations: geometric language, $U : M \rightarrow N$ must be **restricted harmonic**

$$E(U) = E_6(U) + E_0(U) + \varepsilon E_2(U)$$

- $E(U)$ should be stationary for all smooth variations U_t of U
- Choose $U_t = U \circ \psi_t$, ψ_t a curve through Id in $S\text{Diff}(M)$
- E_6 and E_0 are $S\text{Diff}$ invariant! So E_2 must be stationary for these variations: geometric language, $U : M \rightarrow N$ must be **restricted harmonic**
- Strain tensor $\mathcal{D}_{ij} = -\frac{1}{2} \text{tr}(L_i L_j)$, or better $\mathcal{D} = \mathcal{D}_{ij} dx_i dx_j$

$$E(U) = E_6(U) + E_0(U) + \varepsilon E_2(U)$$

- $E(U)$ should be stationary for all smooth variations U_t of U
- Choose $U_t = U \circ \psi_t$, ψ_t a curve through Id in $S\text{Diff}(M)$
- E_6 and E_0 are $S\text{Diff}$ invariant! So E_2 must be stationary for these variations: geometric language, $U : M \rightarrow N$ must be **restricted harmonic**
- Strain tensor $\mathcal{D}_{ij} = -\frac{1}{2} \text{tr}(L_i L_j)$, or better $\mathcal{D} = \mathcal{D}_{ij} dx_i dx_j$
- U restricted harmonic iff $\text{div} \mathcal{D}$ is exact

$$\partial_i \partial_k \mathcal{D}_{kj} dx_i \wedge dx_j = 0$$

$$E(U) = E_6(U) + E_0(U) + \varepsilon E_2(U)$$

- $E(U)$ should be stationary for all smooth variations U_t of U
- Choose $U_t = U \circ \psi_t$, ψ_t a curve through Id in $S\text{Diff}(M)$
- E_6 and E_0 are $S\text{Diff}$ invariant! So E_2 must be stationary for these variations: geometric language, $U : M \rightarrow N$ must be **restricted harmonic**
- Strain tensor $\mathcal{D}_{ij} = -\frac{1}{2} \text{tr}(L_i L_j)$, or better $\mathcal{D} = \mathcal{D}_{ij} dx_i dx_j$
- U restricted harmonic iff $\text{div} \mathcal{D}$ is exact

$$\partial_i \partial_k \mathcal{D}_{kj} dx_i \wedge dx_j = 0$$

- True for all $\varepsilon > 0$. So if $U \xrightarrow{\varepsilon \rightarrow 0} U_{BPS}$, this should be RH also
- Bad news: $U_B = U_H \circ \psi_B$ isn't (failure gets worse as B increases)

Harland's unbound model ($E = E_4 + E_0$)

$$E = -\frac{1}{16} \int_{\mathbb{R}^3} \text{tr}([L_i, L_j][L_i, L_j]) + \int_{\mathbb{R}^3} V(U)$$

Harland's unbound model ($E = E_4 + E_0$)

$$E = -\frac{1}{16} \int_{\mathbb{R}^3} \text{tr}([L_i, L_j][L_i, L_j]) + \int_{\mathbb{R}^3} V(U)$$

- Harland's energy bound:

$$E \geq \underbrace{4(2\pi^2)\langle V^{1/4} \rangle}_C B$$

Proof uses AM-GM and Hölder inequalities

Harland's unbound model ($E = E_4 + E_0$)

$$E = -\frac{1}{16} \int_{\mathbb{R}^3} \text{tr}([L_i, L_j][L_i, L_j]) + \int_{\mathbb{R}^3} V(U)$$

- Harland's energy bound:

$$E \geq \underbrace{4(2\pi^2)\langle V^{1/4} \rangle}_C B$$

Proof uses AM-GM and Hölder inequalities

- Crucial step (Manton): rewrite E_4 in terms of eigenvalues of \mathcal{D}

$$E_4 = \int_M (\lambda_1^2 \lambda_2^2 + \lambda_2^2 \lambda_3^2 + \lambda_3^2 \lambda_1^2)$$

Harland's unbound model ($E = E_4 + E_0$)

$$E = -\frac{1}{16} \int_{\mathbb{R}^3} \text{tr}([L_i, L_j][L_i, L_j]) + \int_{\mathbb{R}^3} V(U)$$

- Harland's energy bound:

$$E \geq \underbrace{4(2\pi^2)\langle V^{1/4} \rangle}_C B$$

Proof uses AM-GM and Hölder inequalities

- Crucial step (Manton): rewrite E_4 in terms of eigenvalues of \mathcal{D}

$$E_4 = \int_M (\lambda_1^2 \lambda_2^2 + \lambda_2^2 \lambda_3^2 + \lambda_3^2 \lambda_1^2)$$

- Bound attained iff

$$\lambda_1^2 = \lambda_2^2 = \lambda_3^2 = V^{1/2}$$

everywhere: $U : \mathbb{R}^3 \rightarrow S^3$ must be conformal with conformal factor $\sqrt{V(U)}$

Harland's unbound model ($E = E_4 + E_0$)

- Essentially unique solution: $U : \mathbb{R}^3 \rightarrow S^3$ is inverse stereographic projection, and

$$V(U) = V_{quartic}(U) = \left(\frac{1}{2} \operatorname{tr}(\mathbb{I}_2 - U) \right)^4$$

- Bound only saturated for $B = 1$. For $B \geq 2$, $E(U) > CB$, so model is **unbound**

$$E_\varepsilon(U) = E_4 + (1 - \varepsilon)E_0^{quartic} + \varepsilon(E_2 + E_0^{pion})$$

- $\varepsilon = 0$ Harland's unbound model, $\varepsilon = 1$ usual model with massive pions

$$E_\varepsilon(U) = E_4 + (1 - \varepsilon)E_0^{quartic} + \varepsilon(E_2 + E_0^{pion})$$

- $\varepsilon = 0$ Harland's unbound model, $\varepsilon = 1$ usual model with massive pions
- Numerics: minimize using conjugate gradient method, start at $\varepsilon = 1$, reduce ε
- Get "realistic" binding energies for $\varepsilon \approx 0.05$, easily accessible to numerics

$$E_\varepsilon(U) = E_4 + (1 - \varepsilon)E_0^{quartic} + \varepsilon(E_2 + E_0^{pion})$$


- $\varepsilon = 0$ Harland's unbound model, $\varepsilon = 1$ usual model with massive pions
- Numerics: minimize using conjugate gradient method, start at $\varepsilon = 1$, reduce ε
- Get "realistic" binding energies for $\varepsilon \approx 0.05$, easily accessible to numerics
- Skyrmions are **lightly bound**: $B = 1$ units occupying subsets of FCC lattice in maximally attractive internal orientation

$$E_\varepsilon(U) = E_4 + (1 - \varepsilon)E_0^{\text{quartic}} + \varepsilon(E_2 + E_0^{\text{pion}})$$


- $\varepsilon = 0$ Harland's unbound model, $\varepsilon = 1$ usual model with massive pions
- Numerics: minimize using conjugate gradient method, start at $\varepsilon = 1$, reduce ε
- Get “realistic” binding energies for $\varepsilon \approx 0.05$, easily accessible to numerics
- Skyrmions are **lightly bound**: $B = 1$ units occupying subsets of FCC lattice in maximally attractive internal orientation
- Many nearly degenerate local minima
- Minima tend to have much less symmetry than in usual $E_2 + E_4$ model

Lightly bound skyrmions


$$\mathcal{B} = 0.1\mathcal{B}_{max}$$



Lightly bound skyrmions



Classical binding energies: summary



Point skyrmion model

- General unit skyrmion

$$U(\mathbf{x}) = U_H(R(\mathbf{x} - \mathbf{x}_0))$$

position \mathbf{x}_0 , orientation $R \in SO(3)$

- Interaction energy of Skyrmion pair at (\mathbf{x}_1, R_1) , (\mathbf{x}_2, R_2) depends only on $\mathbf{X} = \mathbf{x}_1 - \mathbf{x}_2$ and $R = R_1^{-1}R_2$
- Assumption/approximation

$$V_{int} = V_0(|\mathbf{X}|) + V_1(|\mathbf{X}|) \text{tr} R + V_2(|\mathbf{X}|) \frac{\mathbf{X} \cdot R\mathbf{X}}{|\mathbf{X}|^2}$$

- Find V_0, V_1, V_2 by fitting to classical scattering solutions

Point skyrmion model

- General unit skyrmion

$$U(\mathbf{x}) = U_H(R(\mathbf{x} - \mathbf{x}_0))$$

position \mathbf{x}_0 , orientation $R \in SO(3)$

- Interaction energy of Skyrmion pair at (\mathbf{x}_1, R_1) , (\mathbf{x}_2, R_2) depends only on $\mathbf{X} = \mathbf{x}_1 - \mathbf{x}_2$ and $R = R_1^{-1}R_2$
- Assumption/approximation

$$V_{int} = V_0(|\mathbf{X}|) + V_1(|\mathbf{X}|) \text{tr} R + V_2(|\mathbf{X}|) \frac{\mathbf{X} \cdot R\mathbf{X}}{|\mathbf{X}|^2}$$

- Find V_0, V_1, V_2 by fitting to classical scattering solutions
- Very simple point particle approximation to Skyrme energy

$$E_{pp}(\mathbf{x}_1, \dots, \mathbf{x}_B, R_1, \dots, R_B) = BE(U_H) + \sum_{1 \leq a < b \leq B} V_{int}(|\mathbf{x}_a - \mathbf{x}_b|, R_a^{-1}R_b)$$

- Does remarkably well: for $1 \leq B \leq 8$ reproduces all local minima, with correct energy ordering **except** reverses **6a** and **6b**

Point skyrmion model (H+G+JMS+Maybe+Kirk)




3



4



5(a)



5(b)*



6(a)



6(b)



6(c)*



7(a)*



7(b)



8(a)



8(b)



8(c)

- Let it loose on $9 \leq B \leq 23$
- Can automate rigid body quantization procedure (not entirely trivial)
- Results modest: binding energies get inflated (as usual), spin/isospin predictions often unphysical

Point skyrmion model: rigid body quantization

| Name | Bonds | Colour count | Classical energy | Symmetry group | I | J | Quantum energy | Experiment |
|------|-------|--------------|------------------|----------------|-----|-----|----------------|-------------------|
| 2a | 1 | 1,1,0,0 | -0.310 | D_2 | 0 | 1 | 3.813 | $^2\text{H}_1$ |
| 3a | 3 | 1,1,1,0 | -0.931 | C_3 | 1/2 | 1/2 | 1.106 | $^3\text{He}_2$ |
| 4a | 6 | 1,1,1,1 | -1.862 | T | 0 | 0 | -1.862 | $^4\text{He}_2$ |
| 5a | 8 | 2,1,1,1 | -2.338 | 1 | 1/2 | 1/2 | -1.167 | |
| 5b | 8 | 2,2,1,0 | -2.185 | C_4 | 1/2 | 3/2 | -0.700 | $^5\text{He}_2$ |
| 6a | 12 | 2,2,2,0 | -3.229 | O | 2 | 1 | 4.275 | |
| 6b | 11* | 2,2,1,1 | -3.117 | D_2 | 0 | 1 | -2.973 | $^6\text{Li}_3$ |
| 6c | 11* | 2,2,1,1 | -3.046 | 1 | 0 | 0 | -3.046 | |
| 7a | 15 | 2,2,2,1 | -4.057 | C_3 | 1/2 | 1/2 | -3.210 | |
| 8a | 18 | 2,2,2,2 | -4.889 | D_3 | 0 | 0 | -4.889 | $^8\text{Be}_4$ |
| 8b | 18 | 2,2,2,2 | -4.869 | C_2 | 0 | 1 | -4.769 | |
| 9a | 21 | 3,2,2,2 | -5.664 | C_3 | 1/2 | 1/2 | -5.024 | |
| 9b | 21 | 3,2,2,2 | -5.598 | 1 | 1/2 | 1/2 | -4.956 | |
| 10a | 25 | 3,3,2,2 | -6.443 | D_2 | 0 | 1 | -6.352 | |
| 10b | 24* | 4,2,2,2 | -6.442 | T | 0 | 0 | -6.442 | |
| 11a | 28 | 3,3,3,2 | -7.261 | 1 | 1/2 | 1/2 | -6.736 | |
| 12a | 31* | 3,3,3,3 | -8.081 | C_2 | 0 | 0 | -8.081 | $^{12}\text{C}_6$ |
| 12b | 32 | 3,3,3,3 | -8.066 | 1 | 0 | 0 | -8.066 | |
| 13a | 36 | 4,3,3,3 | -9.016 | C_3 | 1/2 | 1/2 | -8.575 | $^{13}\text{C}_6$ |
| 14a | 39* | 4,4,3,3 | -9.821 | 1 | 0 | 0 | -9.821 | |
| 15a | 43* | 4,4,4,3 | -10.653 | 1 | 1/2 | 1/2 | -10.272 | $^{15}\text{N}_7$ |
| 15b | 42** | 4,4,4,3 | -10.627 | 1 | 1/2 | 1/2 | -10.247 | $^{15}\text{N}_7$ |
| 15c | 43* | 4,4,4,3 | -10.584 | 1 | 1/2 | 1/2 | -10.202 | $^{15}\text{N}_7$ |
| 16a | 48 | 4,4,4,4 | -11.771 | T | 0 | 0 | -11.771 | $^{16}\text{O}_8$ |
| 17a | 51* | 5,4,4,4 | -12.563 | C_3 | 1/2 | 1/2 | -12.228 | |
| 18a | 54** | 5,5,4,4 | -13.356 | C_2 | 0 | 0 | -13.356 | |
| 18b | 56 | 6,4,4,4 | -13.340 | C_4 | 0 | 0 | -13.340 | |
| 19a | 60 | 5,5,5,4 | -14.251 | C_3 | 1/2 | 1/2 | -13.951 | $^{19}\text{F}_9$ |
| 19b | 60 | 7,4,4,4 | -14.244 | O | 1/2 | 1/2 | -13.946 | $^{19}\text{F}_9$ |
| 19c | 58** | 5,5,5,4 | -14.178 | 1 | 1/2 | 1/2 | -13.879 | $^{19}\text{F}_9$ |
| 19d | 59* | 5,5,5,4 | -14.164 | 1 | 1/2 | 1/2 | -13.864 | $^{19}\text{F}_9$ |

- Near BPS model ($E_6 + E_0 + \varepsilon E_2$)
 - Skyrmions seem to keep conventional symmetries
 - Has (approx) $SDiff$ invariance: liquid drop model
 - Need $\varepsilon \approx 0.014$ to get realistic B.E.s, much too small for reliable numerics
 - No idea what limit BPS skyrmions actually are. $U_H \circ \psi_B$ certainly wrong.

- Near BPS model ($E_6 + E_0 + \varepsilon E_2$)
 - Skyrmions seem to keep conventional symmetries
 - Has (approx) *SDiff* invariance: liquid drop model
 - Need $\varepsilon \approx 0.014$ to get realistic B.E.s, much too small for reliable numerics
 - No idea what limit BPS skyrmions actually are. $U_H \circ \psi_B$ certainly wrong.
- Lightly bound model ($E_4 + E_0 + \varepsilon(E_2 + E_0^\pi - E_0)$)
 - Numerically tractable at very low ε
 - $\varepsilon \approx 0.05$ yields realistic B.E.s
 - Skyrmions resemble molecules, subsets of FCC lattice
 - Lose symmetries, many nearly degenerate minima
 - Simple and reliable point particle model
 - Has inspired new initial data for conventional model at high B (Manton et al)
 - Good laboratory for more advanced quantization techniques

Summary: other approaches

- **Loosely** bound model (Gudnason): $E = E_4 + E_0 + \varepsilon(E_2 + E_0^\pi)$
but with

$$E_0 = \int_M [\text{tr}(\mathbb{I}_2 - U)]^2 \quad \text{instead of} \quad E_0 = \int_M [\text{tr}(\mathbb{I}_2 - U)]^4.$$

Gets low classical B.E. without losing as much symmetry as lightly bound model.

Summary: other approaches

- **Loosely** bound model (Gudnason): $E = E_4 + E_0 + \varepsilon(E_2 + E_0^\pi)$
but with

$$E_0 = \int_M [\text{tr}(\mathbb{I}_2 - U)]^2 \quad \text{instead of} \quad E_0 = \int_M [\text{tr}(\mathbb{I}_2 - U)]^4.$$

Gets low classical B.E. without losing as much symmetry as lightly bound model.

- Holography (Sutcliffe):
 - Interpret pure YM on M^4 as Skyrme model (on \mathbb{R}^3) coupled to infinite tower of vector mesons
 - Get near BPS theory by truncating meson tower
 - $N = 1$ modest reduction in B.E.s
 - $N = 2$ a lot better (Sutcliffe, Naya)
 - Price: extremely complicated numerical problem
 - Advantage: vector meson coupling interesting for other reasons