

[QMS2020 invited talk]

## **Visualization of the inhomogeneous superfluid in an iron-based superconductor**

Doohee Cho<sup>1</sup>, Koen M Bastiaans<sup>2</sup>, Damianos Chatzopoulos<sup>2</sup>, Gen-Da Gu<sup>2,3</sup>, Milan P Allan<sup>2</sup>

<sup>1</sup> *Department of Physics, Yonsei University, Seoul, Republic of Korea*

<sup>2</sup> *Leiden Institute of Physics, Leiden University, Leiden, the Netherlands*

<sup>3</sup> *Condensed Matter Physics and Materials Science Department, Brookhaven National Laboratory, Upton, NY, USA*

email: dooheecho@yonsei.ac.kr

Cooper pairs can tunnel through the vacuum barrier between two superconducting electrodes, known as a Josephson junction. One can extract the superconducting order parameter from the current-voltage curves close to zero bias. We use atomic-resolution Josephson scanning tunneling microscopy (STM) to visualize the spatial variations of the superfluid in the iron-based superconductor FeTe<sub>0.55</sub>Se<sub>0.45</sub>. To measure Cooper-pair tunneling, a Pb-coated tip was employed and its superconductivity was thoroughly verified by measuring noise enhancement induced by Andreev-reflection on Pb(111) surface. By simultaneously acquiring the topographic and electronic properties, we find that this inhomogeneity in the superfluid is not caused by structural disorder or strong inter-pocket scattering, and does not correlate with variations in the energy of the Cooper pair-breaking gap. Instead, we see a clear spatial correlation between superfluid density and the quasiparticle strength, defined as the height of the coherence peak, on a local scale. Our results shed light on the interplay between superconductivity and quasiparticle character that has been observed by photoemission experiments across the critical temperature in the unconventional superconductors.